

Fax: 1-832-243-6017



A Reliable Research Partner in Life Science and Medicine

## FITC Anti-Mouse CD23 Antibody[B3B4]

Catalog No.E-AB-F1178UCReactivityMouseStorageStore at 2~8°C, Avoid freeze / thaw cyclesApplicationsFCM

**Important Note:** Centrifuge before opening to ensure complete recovery of vial contents.

## **Antigen Information**

Alternate Names Fcer2,Fcer2a,Lymphocyte IgE receptor,Fc-epsilon-RII

Uniprot ID P20693

Background CD23 is a 45 kD protein also known as low affinity IgE Fc receptor, Fc&RII, BLAST-2, Ly-42, or

B6. It is a member of the Ig family, expressed on conventional B (but not B-1) cells and follicular

dendritic cells. CD23 responds to high levels of IgE by downregulating IgE secretion.

#### **Product Details**

 $\begin{tabular}{lll} Form & Liquid \\ Concentration & 0.5 mg/mL \\ Size & 25 \mu g/100 \mu g \\ Clone No. & B3B4 \\ Host & Rat \\ \end{tabular}$ 

 $\begin{tabular}{lll} \textbf{Isotype} & Rat IgG2a, \kappa \\ \textbf{Reactivity} & Mouse \\ \textbf{Application} & FCM \\ \end{tabular}$ 

**Isotype Control** FITC Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09833C]

**Storage Buffer** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

**Shipping** Biological ice pack at 4 °C **Stability & Storage** Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light.Do not freeze.

This product is guaranteed up to one year from purchase.

For Research Use Only

Web: www.elabscience.com Email: techsupport@elabscience.com

A Reliable Research Partner in Life Science and Medicine

# **Fluorophore**


Conjugation: FITC

FITC is designed to be excited by the Blue laser (488 nm) and detected using an optical filter centered near 530 nm (e.g., a 525/40 nm bandpass filter).

## **Recommended usage**

Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1  $\mu g/10^6$  cells in  $100~\mu L$  volume].

### **Product data**



C57BL/6 murine splenocytes are stained with APC Anti-Mouse CD45R/B220 Antibody and FITC Anti-Mouse CD23 Antibody[B3B4] (Left). Splenocytes are stained with APC Anti-Mouse CD45R/B220 Antibody and FITC Rat IgG2a,  $\kappa$  Isotype Control[2A3] (Right).

### **Related Information**

- 1. Sample Preparation for Flow Cytometry <a href="https://www.elabscience.com/List-detail-5594.html">https://www.elabscience.com/List-detail-5594.html</a>
- 2. Staining Cell Surface Targets for Flow Cytometry <a href="https://www.elabscience.com/List-detail-5568.html">https://www.elabscience.com/List-detail-5568.html</a>
- 3. Flow Cytometry Troubleshooting Tips <a href="https://www.elabscience.com/List-detail-5593.html">https://www.elabscience.com/List-detail-5593.html</a>
- 4. How to select the appropriate detection channel through the spectrogram? <a href="https://www.elabscience.com/List-detail-459742.html">https://www.elabscience.com/List-detail-459742.html</a>

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086

 Web: <a href="www.elabscience.com">www.elabscience.com</a>
 Email: <a href="techsupport@elabscience.com">techsupport@elabscience.com</a>