# Elab Fluor® 647 Anti-Human CD197/CCR7 Antibody[G043H7]

| | | | |
|---|---|---|---|
| **Catalog No.** | E-AB-F1159M | **Reactivity** | Human |
| **Storage** | Store at 2~8℃, Avoid freeze / thaw cycles | **Applications** | FCM |

**Important Note:** Centrifuge before opening to ensure complete recovery of vial contents.

## Antigen Information

| | |
|---|---|
| **Alternate Names** | CMKBR7, EBI1, EVI1,CCR-7,CDw197 |
| **Uniprot ID** | P32248 |
| **Background** | CCR7, also known as CD197, is a chemokine receptor that binds CCL19 and CCL21. CCR7 and its ligands link innate and adaptive immunity by affecting interactions between T cells and dendritic cells and their downstream effect. Naïve T cells enter the lymph node through high endothelial venules, which express CCL21. Dendritic cells and macrophages enter the lymph node through afferent lymphatics. The encounter of T cells and dendritic cells in the T cell zone is CCR7-dependent. In addition, during immunological surveillance, B cells recirculate between B-cell-rich compartments (follicles or B cell zones) in secondary lymphoid organs, surveying for antigen. After antigen binding, B cells move to the boundary of B and T zones to interact with T-helper cells; this B cell migration is directed by CCR7 and its ligands. CCR7-positive cancer cell expression has been associated with lymph node metastasis. |

## Product Details

| | |
|---|---|
| **Form** | Liquid |
| **Size** | 20Tests/100Tests/100Tests×2 |
| **Clone No.** | G043H7 |
| **Host** | Mouse |
| **Isotype** | Mouse IgG2a, κ |
| **Reactivity** | Human |
| **Application** | FCM |
| **Isotype Control** | Elab Fluor® 647 Mouse IgG2a, κ Isotype Control[C1.18.4] [Product E-AB-F09802M] |
| **Storage Buffer** | Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant. |
| **Shipping** | Biological ice pack at 4 ℃ |
| **Stability & Storage** | Keep as concentrated solution.<br>Store at 2~8℃ and protected from prolonged exposure to light.Do not freeze.<br>This product is guaranteed up to one year from purchase. |

## Fluorophore

**Conjugation:** Elab Fluor® 647

Elab Fluor® 647 is designed to be excited by the Red laser (627-640 nm) and detected using an optical filter centered near 670 nm (e.g., a 660/20 nm bandpass filter).

## Recommended usage

Each lot of this antibody is quality control tested by flow cytometric analysis. **The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood).** Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

## Product data




Human peripheral blood lymphocytes are stained with Elab Fluor® 647 Anti-Human CD197/CCR7 Antibody and FITC Anti-Human CD3 Antibody (Left). Lymphocytes stained with FITC Anti-Human CD3 Antibody and Elab Fluor® 647 Mouse IgG2a, κ Isotype Control (Right) are used as control.

## Related Information

1. Sample Preparation for Flow Cytometry https://www.elabscience.com/List-detail-5594.html
2. Staining Cell Surface Targets for Flow Cytometry https://www.elabscience.com/List-detail-5568.html
3. Flow Cytometry Troubleshooting Tips https://www.elabscience.com/List-detail-5593.html
4. How to select the appropriate detection channel through the spectrogram? https://www.elabscience.com/List-detail-459742.html