



A Reliable Research Partner in Life Science and Medicine

# PE/Cyanine5 Anti-Mouse IgM Antibody[RMM-1]

E-AB-F1190UG Catalog No. Reactivity Mouse Storage Store at 2~8°C, Avoid freeze / thaw cycles **Applications FCM** 

**Important Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### **Antigen Information**

**Alternate Names** Immunoglobulin heavy constant mu,IGHM,Immunoglobulin M

**Uniprot ID** P01872

**Background** IgM is the first immunoglobulin made by B cells in the immune response. Surface IgM is

expressed on the majority of mature B cells.

#### **Product Details**

**Form** Liquid Concentration 0.2 mg/mL  $25\mu g/100\mu g$ Size RMM-1 Clone No. Host Rat

**Isotype** Rat IgG2a, κ Reactivity Mouse **Application FCM** 

PE/Cyanine5 Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09833G] **Isotype Control** 

**Storage Buffer** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

**Shipping** Biological ice pack at 4 °C Stability & Storage Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light.Do not freeze.

This product is guaranteed up to one year from purchase.

For Research Use Only

Fax: 1-832-243-6017 Web: www.elabscience.com Email: techsupport@elabscience.com

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086

A Reliable Research Partner in Life Science and Medicine

## **Fluorophore**


Conjugation: PE/Cyanine5

PE/Cyanine5 is designed to be excited by the Blue (488 nm), Green (532 nm) and yellow-green (561 nm) lasers and detected using an optical filter centered near 670 nm (e.g., a 690/50 nm bandpass filter).

## **Recommended usage**

Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is  $0.1-1~\mu g/10^6$  cells in  $100~\mu L$  volume].

#### **Product data**



C57BL/6 murine splenocytes are stained with PE/Cyanine5 Anti-Mouse IgM Antibody[RMM-1] (filled gray histogram) or PE/Cyanine5 Rat IgG2a, κ Isotype Control (empty black histogram).

#### **Related Information**

- 1. Sample Preparation for Flow Cytometry <a href="https://www.elabscience.com/List-detail-5594.html">https://www.elabscience.com/List-detail-5594.html</a>
- 2. Staining Cell Surface Targets for Flow Cytometry <a href="https://www.elabscience.com/List-detail-5568.html">https://www.elabscience.com/List-detail-5568.html</a>
- 3. Flow Cytometry Troubleshooting Tips <a href="https://www.elabscience.com/List-detail-5593.html">https://www.elabscience.com/List-detail-5593.html</a>
- 4. How to select the appropriate detection channel through the spectrogram? <a href="https://www.elabscience.com/List-detail-459742.html">https://www.elabscience.com/List-detail-459742.html</a>

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086

 Web: <a href="www.elabscience.com">www.elabscience.com</a>
 Email: <a href="techsupport@elabscience.com">techsupport@elabscience.com</a>