



A Reliable Research Partner in Life Science and Medicine

# Elab Fluor® 488 Anti-Mouse H-2 Antibody[M1/42]

E-AB-F1216L Catalog No. Reactivity Mouse Storage Store at 2~8°C, Avoid freeze / thaw cycles **Applications FCM** 

**Important Note:** Centrifuge before opening to ensure complete recovery of vial contents.

## **Antigen Information**

**Alternate Names** Mouse major histocompatibility complex (MHC) H-2, MHC I

**Background** The M1/42 antibody reacts with the H-2 MHC class I alloantigens expressed on nucleated cells

from mice of the a, b, d, j, k, s, and u haplotypes (Stallcup, KC et al, 1981). MHC class I is

involved in antigen presentation to T cells expressing CD3/TCR and CD8 proteins.

#### **Product Details**

**Form** Liquid

50Tests/100Tests/100Tests×2 Size

Clone No. M1/42 Host Rat

Rat IgG2a, κ **Isotype** Reactivity Mouse **Application FCM** 

Elab Fluor<sup>®</sup> 488 Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09832L] **Isotype Control** 

Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant. **Storage Buffer** 

Shipping Biological ice pack at 4 °C Stability & Storage Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light.Do not freeze.

This product is guaranteed up to one year from purchase.

For Research Use Only

Toll-free: 1-888-852-8623 Web: www.elabscience.com Email: techsupport@elabscience.com

Tel: 1-832-243-6086 Fax: 1-832-243-6017

#### **Elabscience Bionovation Inc.**

Fax: 1-832-243-6017



A Reliable Research Partner in Life Science and Medicine

## **Fluorophore**

Conjugation: Elab Fluor® 488

Elab Fluor<sup>®</sup> 488 is designed to be excited by the Blue laser (488 nm) and detected using an optical filter centered near 520 nm (e.g., a 525/40 nm bandpass filter).

### Recommended usage

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

#### **Related Information**

- 1. Sample Preparation for Flow Cytometry <a href="https://www.elabscience.com/List-detail-5594.html">https://www.elabscience.com/List-detail-5594.html</a>
- 2. Staining Cell Surface Targets for Flow Cytometry <a href="https://www.elabscience.com/List-detail-5568.html">https://www.elabscience.com/List-detail-5568.html</a>
- 3. Flow Cytometry Troubleshooting Tips <a href="https://www.elabscience.com/List-detail-5593.html">https://www.elabscience.com/List-detail-5593.html</a>
- 4. How to select the appropriate detection channel through the spectrogram? <a href="https://www.elabscience.com/List-detail-459742.html">https://www.elabscience.com/List-detail-459742.html</a>

For Research Use Only

Web: www.elabscience.com Email: techsupport@elabscience.com

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086