



A Reliable Research Partner in Life Science and Medicine

# PerCP/Cyanine5.5 Anti-Mouse H-2 Antibody[M1/42]

Catalog No. E-AB-F1216UJ Reactivity Mouse Storage Store at 2~8°C, Avoid freeze / thaw cycles **Applications FCM** 

**Important Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### **Antigen Information**

**Alternate Names** Mouse major histocompatibility complex (MHC) H-2, MHC I

**Background** The M1/42 antibody reacts with the H-2 MHC class I alloantigens expressed on nucleated cells

from mice of the a, b, d, j, k, s, and u haplotypes (Stallcup, KC et al, 1981). MHC class I is

involved in antigen presentation to T cells expressing CD3/TCR and CD8 proteins.

#### **Product Details**

**Form** Liquid Concentration 0.2 mg/mL Size  $25 \mu g / 100 \mu g$ M1/42 Clone No. Host Rat

**Isotype** Rat IgG2a, κ Reactivity Mouse **Application FCM** 

PerCP/Cyanine 5.5 Rat IgG2a, κ Isotype Control [2A3] [Product E-AB-F09833J] **Isotype Control** 

**Storage Buffer** Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.

**Shipping** Biological ice pack at 4 °C Stability & Storage Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light.Do not freeze.

This product is guaranteed up to one year from purchase.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com



Fax: 1-832-243-6017



A Reliable Research Partner in Life Science and Medicine

### **Fluorophore**

Conjugation: PerCP/Cyanine5.5

PerCP/Cyanine5.5 is designed to be excited by the blue laser (488 nm) and detected using an optical filter centered near 675 nm (e.g., a 690/50 nm bandpass filter).

## **Recommended usage**

Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1  $\mu g/10^6$  cells in  $100~\mu L$  volume].

#### **Related Information**

- 1. Sample Preparation for Flow Cytometry <a href="https://www.elabscience.com/List-detail-5594.html">https://www.elabscience.com/List-detail-5594.html</a>
- 2. Staining Cell Surface Targets for Flow Cytometry <a href="https://www.elabscience.com/List-detail-5568.html">https://www.elabscience.com/List-detail-5568.html</a>
- 3. Flow Cytometry Troubleshooting Tips <a href="https://www.elabscience.com/List-detail-5593.html">https://www.elabscience.com/List-detail-5593.html</a>
- 4. How to select the appropriate detection channel through the spectrogram? <a href="https://www.elabscience.com/List-detail-459742.html">https://www.elabscience.com/List-detail-459742.html</a>

For Research Use Only

Web: <u>www.elabscience.com</u> Email: <u>techsupport@elabscience.com</u>