

Fax: 1-832-243-6017



A Reliable Research Partner in Life Science and Medicine

AF/LE Purified Anti-Mouse CD132 Antibody[3E12]

Catalog No.E-AB-F10150ReactivityMouseStorageStore at 2~8°C, Avoid freeze / thaw cyclesApplicationsFCM

Important Note: Centrifuge before opening to ensure complete recovery of vial contents.

Antigen Information

Alternate Names Cytokine receptor common subunit gamma,Il2rg,Interleukin-2 receptor subunit gamma,IL-2

receptor subunit gamma, IL-2R subunit gamma, IL-2RG, gammaC, p64, CD132

Uniprot ID P34902

Background CD132 is a 64-70 kD type I transmembrane glycoprotein of the Ig superfamily, and is also known

as common γ chain (γ c) or IL-2 receptor γ subunit. It is expressed broadly on T- and B-lymphocytes, NK cells, monocytes, and granulocytes. CD132 is an essential component of cytokine receptors for IL-2, IL-4, IL-7, IL-9, IL-15, and IL-21. Ligand binding induces tyrosine phosphorylation and initiates signaling through a JAK/STAT pathway. CD132 mutation results in

X-linked severe combined immune deficiency (XSCID).

Product Details

 Form
 Liquid

 Concentration
 0.5 mg/mL

 Size
 50μg/500μg/1mg

Clone No. 3E12 Host Rat

IsotypeRat IgG2b, κReactivityMouseApplicationFCM

Isotype ControlAF/LE Purified Rat IgG2b, κ Isotype Control[LTF-2] [Product E-AB-F098430]

Storage Buffer 0.2 µm filtered in PBS, pH 7.2. Azide Free (AF)/Low Endotoxin (LE): Contains no stabilizers or

stabilizers. Endotoxin level is < 2 EU/mg as Determined by LAL gel clotting assay.

Shipping Biological ice pack at 4 °C

Stability & Storage Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light.Do not freeze.

This product is guaranteed up to one year from purchase.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086

Web: www.elabscience.com
Email: techsupport@elabscience.com





A Reliable Research Partner in Life Science and Medicine

Fluorophore

Conjugation: None (Purified antibody-Azide Free/Low endotoxin)

Recommended usage

Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is $\leq 2.0 \,\mu g$ per 10^6 cells in $100 \,\mu L$ volume or $100 \,\mu L$ of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Related Information

- 1. Sample Preparation for Flow Cytometry https://www.elabscience.com/List-detail-5594.html
- 2. Staining Cell Surface Targets for Flow Cytometry https://www.elabscience.com/List-detail-5568.html
- 3. Flow Cytometry Troubleshooting Tips https://www.elabscience.com/List-detail-5593.html
- 4. How to select the appropriate detection channel through the spectrogram? https://www.elabscience.com/Listdetail-459742.html

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web: www.elabscience.com Email: techsupport@elabscience.com

Fax: 1-832-243-6017