

AF/LE Purified Anti-Mouse CD209b Antibody[22D1]

| | | | |
|--------------------|--|---------------------|-------|
| Catalog No. | E-AB-F10220 | Reactivity | Mouse |
| Storage | Store at 2~8°C, Avoid freeze / thaw cycles | Applications | FCM |

Important Note: Centrifuge before opening to ensure complete recovery of vial contents.

Antigen Information

| | |
|------------------------|--|
| Alternate Names | CD209 antigen-like protein B, Cd209b, DC-SIGN-related protein 1, DC-SIGNR1, OtB7, CD209, Cd209b |
| Uniprot ID | Q8CJ91 |
| Background | CD209B, also known as SIGN-R1, is a mouse C-type lectin receptor predominantly expressed on macrophages in the spleen marginal zone and lymph nodes medulla. CD209B is a mouse homolog of human CD209/DC-SIGN and is involved in innate immune response. CD209B mediates the recognition and uptake of pathogen products, such as lipopolysaccharides (LPS), pneumococcal polysaccharides, and dextrans. CD209B has been demonstrated to facilitate the clearance of encapsulated pneumococcus by directly binding to C1q and activating complement through an immunoglobulin independent pathway. |

Product Details

| | |
|--------------------------------|---|
| Form | Liquid |
| Concentration | 0.5 mg/mL |
| Size | 50µg/500µg/1mg |
| Clone No. | 22D1 |
| Host | Armenian Hamster |
| Isotype | Armenian Hamster IgG |
| Reactivity | Mouse |
| Application | FCM |
| Isotype Control | AF/LE Purified Armenian Hamster IgG Isotype Control[PIP] [Product E-AB-F098530] |
| Storage Buffer | 0.2 µm filtered in PBS, pH 7.2. Azide Free (AF)/Low Endotoxin (LE): Contains no stabilizers or stabilizers. Endotoxin level is < 2 EU/mg as Determined by LAL gel clotting assay. |
| Shipping | Biological ice pack at 4 °C |
| Stability & Storage | Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. This product is guaranteed up to one year from purchase. |

For Research Use Only

Fluorophore

Conjugation: None (Purified antibody-Azide Free/Low endotoxin)

Recommended usage

Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is $\leq 0.5 \mu\text{g}$ per 10^6 cells in $100 \mu\text{L}$ volume or $100 \mu\text{L}$ of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Related Information

1. Sample Preparation for Flow Cytometry <https://www.elabscience.com/List-detail-5594.html>
2. Staining Cell Surface Targets for Flow Cytometry <https://www.elabscience.com/List-detail-5568.html>
3. Flow Cytometry Troubleshooting Tips <https://www.elabscience.com/List-detail-5593.html>
4. How to select the appropriate detection channel through the spectrogram? <https://www.elabscience.com/List-detail-459742.html>