

## AF/LE Purified Anti-Mouse CD120b/TNFR2 Antibody[TR75-54.7]

<b>Catalog No.</b>	E-AB-F10350	<b>Reactivity</b>	Mouse
<b>Storage</b>	Store at 2~8°C, Avoid freeze / thaw cycles	<b>Applications</b>	FCM,Block

**Important Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Antigen Information

<b>Alternate Names</b>	Tumor necrosis factor receptor superfamily member 1B,Tnfrsf1b,Tumor necrosis factor receptor 2,TNF-R2,TNF-RII,TNFR-II,p75,p80 TNF-alpha receptor,CD120b
<b>Uniprot ID</b>	P25119
<b>Background</b>	CD120b is a 75 kD type I transmembrane protein, also known as Tumor Necrosis Factor Receptor Type II (TNFRII) or p75. It is expressed on a variety of cells at low levels; the expression is upregulated upon activation. This receptor binds both TNF- $\alpha$ and LT- $\alpha$ (also known as TNF- $\beta$ ). In association with TRAF1 and TRAF2, the receptor crosslinking induced by TNF- $\alpha$ or LT- $\alpha$ trimers is critical for signal transduction, leading to apoptosis, NF- $\kappa$ B activation, increased expression of proinflammatory genes, tumor necrosis, and cell differentiation depending on cell type and differentiation state.

### Product Details

<b>Form</b>	Liquid
<b>Concentration</b>	0.5 mg/mL
<b>Size</b>	50 $\mu$ g/500 $\mu$ g/1mg
<b>Clone No.</b>	TR75-54.7
<b>Host</b>	Armenian Hamster
<b>Isotype</b>	Armenian Hamster IgG
<b>Reactivity</b>	Mouse
<b>Application</b>	FCM,Block
<b>Isotype Control</b>	<a href="#">AF/LE Purified Armenian Hamster IgG Isotype Control[PIP] [Product E-AB-F098530]</a>
<b>Storage Buffer</b>	0.2 $\mu$ m filtered in PBS, pH 7.2. Azide Free (AF)/Low Endotoxin (LE): Contains no stabilizers or stabilizers. Endotoxin level is < 2 EU/mg as Determined by LAL gel clotting assay.
<b>Shipping</b>	Biological ice pack at 4 °C
<b>Stability &amp; Storage</b>	Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light.Do not freeze. This product is guaranteed up to one year from purchase.

### For Research Use Only

## Fluorophore

**Conjugation:** None (Purified antibody-Azide Free/Low endotoxin)

## Recommended usage

Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is  $\leq 0.25 \mu\text{g}$  per  $10^6$  cells in  $100 \mu\text{L}$  volume or  $100 \mu\text{L}$  of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

## Related Information

1. Sample Preparation for Flow Cytometry <https://www.elabscience.com/List-detail-5594.html>
2. Staining Cell Surface Targets for Flow Cytometry <https://www.elabscience.com/List-detail-5568.html>
3. Flow Cytometry Troubleshooting Tips <https://www.elabscience.com/List-detail-5593.html>
4. How to select the appropriate detection channel through the spectrogram? <https://www.elabscience.com/List-detail-459742.html>