## **Elabscience**®

## AF/LE Purified Anti-Mouse CD119 Antibody[GR-20]

Catalog No.E-AB-F11150StorageStore at 2~8°C, Avoid freeze / thaw cycles

ReactivityMouseApplicationsFCM

**Important Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### **Antigen Information**

Alternate Names	Interferon gamma receptor 1, Ifngr1, IFN-gamma-R1, IFN-gamma-R-alpha, CD119
Uniprot ID	P15261
Background	CDw119 is a 90 kD immunoglobulin superfamily member, also known as IFN- $\gamma R\alpha$ chain. It is a class II cytokine receptor family member that serves as a IFN- $\gamma$ -binding chain associated with the IFN- $\gamma$ $\beta$ chain also known as AF-1. In addition to ligand binding, CDw119 participates in ligand
	trafficking. CDw119 is expressed on T and B cells, NK cells, fibroblasts, endothelial, and epithelial cells. Binding of IFN- $\gamma$ induces receptor dimerization, internalization, Jak1 and Jak2 protein kinase activation and, ultimately, STAT1 activation. IFN- $\gamma$ initiates and regulates a variety of immune responses.

#### **Product Details**

Form	Liquid
Concentration	0.5 mg/mL
Size	50μg/500μg/1mg
Clone No.	GR-20
Host	Rat
Isotype	Rat IgG2a, κ
Reactivity	Mouse
Application	FCM
Isotype Control	AF/LE Purified Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F098330]
Storage Buffer	0.2 µm filtered in PBS, pH 7.2. Azide Free (AF)/Low Endotoxin (LE): Contains no stabilizers or
	stabilizers. Endotoxin level is $< 2$ EU/mg as Determined by LAL gel clotting assay.
Shipping	Biological ice pack at 4 °C
Stability & Storage	Keep as concentrated solution.
	Store at 2~8°C and protected from prolonged exposure to light.Do not freeze.
	This product is guaranteed up to one year from purchase.

**For Research Use Only** 

# **Elabscience**®

## Fluorophore

Conjugation: None (Purified antibody-Azide Free/Low endotoxin)

### **Recommended usage**

Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is  $\leq 2.0 \ \mu g$  per 10<sup>6</sup> cells in 100  $\mu L$  volume or 100  $\mu L$  of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

## **Related Information**

- 1. Sample Preparation for Flow Cytometry <u>https://www.elabscience.com/List-detail-5594.html</u>
- 2. Staining Cell Surface Targets for Flow Cytometry https://www.elabscience.com/List-detail-5568.html
- 3. Flow Cytometry Troubleshooting Tips https://www.elabscience.com/List-detail-5593.html
- 4. How to select the appropriate detection channel through the spectrogram? <u>https://www.elabscience.com/List-detail-459742.html</u>

#### For Research Use Only