



A Reliable Research Partner in Life Science and Medicine

Purified Anti-Human CD28 Antibody[CD28.2]

Catalog No. E-AB-F1195A Reactivity Human Storage Store at 2~8°C, Avoid freeze / thaw cycles **Applications FCM**

Important Note: Centrifuge before opening to ensure complete recovery of vial contents.

Antigen Information

Alternate Names PAS2,MNS,HGpMiX,GYPA,GPSAT

Uniprot ID P10747

Background CD28 is a 44 kD disulfide-linked homodimeric type I glycoprotein. It is a member of the

> immunoglobulin superfamily and is also known as T44 or Tp44. CD28 is expressed on most T lineage cells, NK cell subsets, and plasma cells. CD28 binds both CD80 and CD86 using a highly conserved motif MYPPY in the CDR3-like loop. CD28 is considered a major co-stimulatory molecule, inducing T lymphocyte activation and IL-2 synthesis, and preventing cell death. In vitro studies indicate that ligation of CD28 on T cells by CD80 and CD86 on antigen presenting cells

provides a costimulatory signal required for T cell activation and proliferation.

Product Details

Form Liquid Concentration 0.5 mg/mL $25 \mu g / 100 \mu g$ Size Clone No. CD28.2 Host Mouse

Mouse IgG1, κ **Isotype** Reactivity Human **Application FCM**

Isotype Control Purified Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09793A]

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

Shipping Biological ice pack at 4 °C Stability & Storage Keep as concentrated solution. Store at 2~8°C .Do not freeze.

This product is guaranteed up to one year from purchase.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Email: techsupport@elabscience.com Web: www.elabscience.com

Fax: 1-832-243-6017

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

Recommended usage

Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is $\leq 0.5~\mu g$ per 10^6 cells in $100~\mu L$ volume or $100~\mu L$ of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Related Information

- 1. Sample Preparation for Flow Cytometry https://www.elabscience.com/List-detail-5594.html
- 2. Staining Cell Surface Targets for Flow Cytometry https://www.elabscience.com/List-detail-5568.html
- 3. Flow Cytometry Troubleshooting Tips https://www.elabscience.com/List-detail-5593.html
- 4. How to select the appropriate detection channel through the spectrogram? https://www.elabscience.com/List-detail-459742.html

For Research Use Only

Web: www.elabscience.com Email: techsupport@elabscience.com

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017