

Recombinant Human HVEM/TNFRSF14 Protein (Fc Tag)



Catalog Number:PKSH033487

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Tumor Necrosis Factor Receptor Superfamily Member 14;Herpes Virus Entry Mediator A;Herpesvirus Entry Mediator A;HveA;Tumor Necrosis Factor Receptor-Like 2;TR2;CD270
Species	Human
Expression Host	HEK293 Cells
Sequence	Leu39-Val202
Accession	Q92956
Calculated Molecular Weight	44.5 kDa
Observed molecular weight	50-60 kDa
Tag	C-Fc

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Herpesvirus entry mediator (HVEM) is a type I membrane protein in the TNF receptor superfamily; and it can both promote and inhibit T cell activity. HVEM is highly expressed on na[?]ve CD4⁺ T cells; CD8⁺ T memory cells; regulatory T cells; dendritic cells; monocytes; and neutrophils. It functions as a receptor for BTLA; CD160; LIGHT/TNFSF14; and Lymphotoxin-alpha. Ligation of HVEM by LIGHT triggers T cell; monocyte; and neutrophil activation and contributes to Th1 inflammation and cardiac allograft rejection. In contrast; HVEM binding to CD160 or BTLA suppresses T cell and dendritic cell activation and dampens intestinal inflammation. HVEM enhances the development of CD8⁺ T cell memory

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Recombinant Human HVEM/TNFRSF14 Protein (Fc Tag)



Catalog Number:PKSH033487

and Treg function. It is additionally expressed on intestinal epithelial cells; where its binding by intraepithelial lymphocyte (IEL) expressed CD160 promotes epithelial integrity and host defense. The herpesvirus envelope glycoprotein gD; which binds HVEM to initiate membrane fusion; can antagonize both BTLA and LIGHT binding.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Tel: 1-832-243-6086

Fax: 1-832-243-6017

Web: www.elabscience.com

Email: techsupport@elabscience.com