

# Recombinant Human HVEM/TNFRSF14 Protein (Fc Tag)

Catalog Number:PKSH033487



**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

<b>Synonyms</b>	Tumor Necrosis Factor Receptor Superfamily Member 14;Herpes Virus Entry Mediator A;Herpesvirus Entry Mediator A;HveA;Tumor Necrosis Factor Receptor-Like 2;TR2;CD270
<b>Species</b>	Human
<b>Expression Host</b>	HEK293 Cells
<b>Sequence</b>	Leu39-Val202
<b>Accession</b>	Q92956
<b>Calculated Molecular Weight</b>	44.5 kDa
<b>Observed molecular weight</b>	50-60 kDa
<b>Tag</b>	C-Fc

## Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

## Data



> 95 % as determined by reducing SDS-PAGE.

## Background

Herpesvirus entry mediator (HVEM) is a type I membrane protein in the TNF receptor superfamily; and it can both promote and inhibit T cell activity. HVEM is highly expressed on na?ve CD4+ T cells; CD8+ T memory cells; regulatory T cells; dendritic cells; monocytes; and neutrophils. It functions as a receptor for BTLA; CD160; LIGHT/TNFSF14; and Lymphotoxin-alpha. Ligation of HVEM by LIGHT triggers T cell; monocyte; and neutrophil activation and contributes to Th1 inflammation and cardiac allograft rejection. In contrast; HVEM binding to CD160 or BTLA suppresses T cell and dendritic cell activation and dampens intestinal inflammation. HVEM enhances the development of CD8+ T cell memory

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

# Recombinant Human HVEM/TNFRSF14 Protein (Fc Tag)

Catalog Number: PKSH033487



and Treg function. It is additionally expressed on intestinal epithelial cells; where its binding by intraepithelial lymphocyte (IEL) expressed CD160 promotes epithelial integrity and host defense. The herpesvirus envelope glycoprotein gD; which binds HVEM to initiate membrane fusion; can antagonize both BTLA and LIGHT binding.

---

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017