A Reliable Research Partner in Life Science and Medicine

# **Recombinant Human TGFB1/TGF-beta 1 Protein (His Tag)**

Catalog No. PKSH033495

Note: Centrifuge before opening to ensure complete recovery of vial contents.

### **Description**

Synonyms Transforming Growth Factor Beta-1;TGF-Beta-1;Latency-Associated

Peptide;LAP;TGFB1;TGFB;CED;DPD1;TGF-beta 1

Species Human

Expression Host HEK293 Cells

Sequence Leu30-Ser390(Cys33Ser)

Accession P01137

Calculated Molecular Weight 29.3&12.8 kDa
Observed molecular weight 38-55&13 kDa

Tag N-His

**Bioactivity** Not validated for activity

## **Properties**

**Purity** > 95 % as determined by reducing SDS-PAGE.

**Endotoxin** < 1.0 EU per µg of the protein as determined by the LAL method.

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation** Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

#### Data



> 95 % as determined by reducing SDS-PAGE.

# **Background**

Transforming Growth Factor  $\beta$ -1 (TGF $\beta$ -1) is a secreted protein which belongs to the TGF- $\beta$  family. TGF $\beta$ -1 is

### For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com

Email: techsupport@elabscience.com

# **Elabscience Bionovation Inc.**



A Reliable Research Partner in Life Science and Medicine

abundantly expressed in bone; articular cartilage and chondrocytes and is increased in osteoarthritis (OA). TGFβ-1 performs many cellular functions; including the control of cell growth; cell proliferation; cell differentiation and apoptosis. The precursor is cleaved into a latency-associated peptide (LAP) and a mature TGFβ-1 peptide. TGFβ-1 may also form heterodimers with other TGFβ family members. It has been found that TGFβ-1 is frequently upregulated in tumor cells. Mutations in this gene results in Camurati-Engelmann disease.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com

Web: www.elabscience.com