

Recombinant Human FGF-2/FGF basic/FGFb Protein

Catalog No. PKSH033503

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Fibroblast growth factor 2;FGF-2;Basic fibroblast growth factor;bFGF;Heparin-binding growth factor 2;HBGF-2;FGF2;FGFB
Species	Human
Expression Host	E.coli
Sequence	Met1-Ser155
Accession	BAG70135.1
Calculated Molecular Weight	17.2 kDa
Observed molecular weight	17 kDa
Tag	None
Bioactivity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.
Shipping	This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel packs. Upon receipt, store it immediately at < -20°C.
Formulation	Supplied as a 0.2 µm filtered solution of 20mM Tris-HCl, 200mM NaCl, pH 7.5.
Reconstitution	Not Applicable

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Fibroblast growth factors (FGF) are a family of heparin-binding secreted proteins that stimulate cell proliferation and differentiation in a wide variety of tissues. FGFs play important roles in diverse biological functions both in vivo and in vitro; including mitogenesis; cellular migration; differentiation; angiogenesis; and wound healing. Human embryonic stem cell (hESC) cultures require FGF basic (also known as FGF-2 or bFGF) in cell culture media to remain in an undifferentiated and pluripotent state. Thermostable FGF basic was engineered for enhanced stability in culture media;

For Research Use Only

without modification of its biological function. FGF basic is a required component of stem cell culture media for maintaining cells in an undifferentiated state. Because FGF basic is unstable; daily media changes are needed. The thermostable FGF basic that supports a 2-day media change schedule; so no media changes are required over a weekend. This thermostable FGF basic was more stable than FGF basic in biochemical studies; and maintained cell growth; pluripotency and differentiation potential with a 2-day feeding schedule.