

# Recombinant Human S100A11 Protein

Catalog Number:PKSH033550



**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

<b>Synonyms</b>	Protein S100-A11;Calgizzarin;Metastatic lymph node gene 70 protein;MLN 70;Protein S100-C;S100 calcium-binding protein A11;S100A11;MLN70;S100C
<b>Species</b>	Human
<b>Expression Host</b>	E.coli
<b>Sequence</b>	Met1-Thr105
<b>Accession</b>	P31949
<b>Calculated Molecular Weight</b>	11.7 kDa
<b>Observed molecular weight</b>	12 kDa
<b>Tag</b>	None

## Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of PBS, 1mM DTT, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

## Data



> 95 % as determined by reducing SDS-PAGE.

## Background

S100A11 is a member of the S100 family of calcium binding proteins. Human S100A11 contains two EF hand motifs and shares 82% amino acid sequence identity with mouse and rat S100A11. It forms covalent homodimers upon transglutamination and also disulfide-linked tetramers. S100A11 is secreted by keratinocytes and can be crosslinked into the cornified envelope of the skin. Dimerization enhances its ability to signal through RAGE on keratinocytes, induce the production of EGF family proteins, and induce cell proliferation. Dimerization also enables S100A11 to bind RAGE on chondrocytes, leading to chondrocyte hypertrophy and catabolism of the cartilage matrix. S100A11 is additionally found in the cytosol where it becomes phosphorylated and translocates to the nucleus in response to DNA damage, RELM alpha

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

# Recombinant Human S100A11 Protein

Catalog Number:PKSH033550



exposure, or elevated extracellular calcium concentrations. Calcium also promotes S100A11 association with S100B as well as Annexins A1, A2, and A6. S100A11-Annexin A2 complexes are recruited to sites of plasma membrane damage where they facilitate membrane repair in migrating cancer cells. S100A11 is upregulated in various cancers and supports tumor cell proliferation, invasion, and migration. In addition, S100A11 is produced in the ovary, and it acts on cumulus cells to inhibit oocyte fertilization.

---

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017