A Reliable Research Partner in Life Science and Medicine

Recombinant Human LFA-3/CD58 Protein (Fc Tag)

Catalog No. PKSH033594

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms Lymphocyte Function-Associated Antigen 3;Surface Glycoprotein

LFA-3;CD58;LFA3;Ag3;CD58 antigen

Species Human

Expression Host

Sequence

Phe29-Arg215

Accession

Calculated Molecular Weight

Observed molecular weight

Tag

HEK293 Cells

Phe29-Arg215

AAH05930

48.6 kDa

60-90 kDa

C-Fc

Bioactivity Not validated for activity

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per μg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Lymphocyte function-associated antigen 3 (LFA-3/CD58) is a single-pass type I membrane protein. CD58 is widely

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: <u>www.elabscience.com</u> Email: <u>techsupport@elabscience.com</u>

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

expressed on hematopoietic and non-hematopoietic human tissue and has been found on leukocytes; erythrocytes; endothelial cells; epithelial cells and fibroblasts of human origin. It is a Ligand of the T-lymphocyte CD2 glycoprotein. This interaction is important in mediating thymocyte interactions with thymic epithelial cells; antigen-independent and -dependent interactions of T-lymphocytes with target cells and antigen-presenting cells and the T-lymphocyte rosetting with erythrocytes. In addition; the LFA-3/CD2 interaction may prime response by both the CD2+ and LFA-3+ cells.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com

Web: www.elabscience.com