

## Recombinant Human CD16b/FCGR3B Protein (aa 17-208, His Tag)

**Catalog No.** PKSH033599

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

|                                    |  |
|------------------------------------|--|
| <b>Synonyms</b>                    | Low affinity immunoglobulin gamma Fc region receptor III-B;Fc-gamma RIII-beta;FcR-10;IgG Fc receptor III-1;FCG3;FCGR3;CD16b and FCGR5B;FCRIII;FCRIIb |
| <b>Species</b>                     | Human  |
| <b>Expression Host</b>             | HEK293 Cells   |
| <b>Sequence</b>                    | Gly17-Gln208   |
| <b>Accession</b>                   | O75015   |
| <b>Calculated Molecular Weight</b> | 22.7 kDa   |
| <b>Observed molecular weight</b>   | 35-65 kDa  |
| <b>Tag</b>                         | C-His  |
| <b>Bioactivity</b>                 | Not validated for activity   |

### Properties

|                       |   |
|-----------------------|---|
| <b>Purity</b>         | > 95 % as determined by reducing SDS-PAGE.  |
| <b>Endotoxin</b>      | < 1.0 EU per µg of the protein as determined by the LAL method.   |
| <b>Storage</b>        | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.             |
| <b>Shipping</b>       | This product is provided as lyophilized powder which is shipped with ice packs.   |
| <b>Formulation</b>    | Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.<br>Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.<br>Please refer to the specific buffer information in the printed manual. |
| <b>Reconstitution</b> | Please refer to the printed manual for detailed information.  |

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

#### For Research Use Only

Low affinity immunoglobulin gamma Fc region receptor III-B; also known as Fc-gamma RIII-beta; FcR-10; IgG Fc receptor III-1; FCG3; FCGR3; CD16b and FCGR3B. FCGR3B is a GPI-anchor membrane protein and contains two Ig-like C2 type domains. FCGR3B can be expressed in orphonuclear leukocytes and stimulated eosinophils. FCGR3B can interact with INPP5D/SHIP1. FCGR3B localizes in the FCGR gene cluster is a CN polymorphic gene involved in the recruitment of polymorphonuclear neutrophils to sites of inflammation and their activation. FCGR3B may serve as a trap for immune complexes in the peripheral circulation which does not activate neutrophils.