A Reliable Research Partner in Life Science and Medicine

Recombinant Human Interleukin-11/IL-11 Protein

Catalog No. PKSH033631

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms Interleukin-11;IL-11;Adipogenesis Inhibitory Factor;AGIF;Oprelvekin;IL11

SpeciesHumanExpression HostP.pastorisSequenceGly23-Leu199

AccessionP20809Calculated Molecular Weight19.0 kDaObserved molecular weight19 kDaTagNone

Bioactivity Not validated for activity

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from a 0.2 µm filtered solution of 20mM PB, 2% Glycine, pH 7.2.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Interleukin 11 (IL-11) is a member of a family of human growth factors that includes human growth hormone, granulocyte colony-stimulating factor, and other growth factors. IL-11 is a thrombopoietic growth factor that directly

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com

Email: techsupport@elabscience.com

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

stimulates the proliferation of hematopoietic stem cells and megakaryocyte progenitor cells and induces megakaryocyte maturation resulting in increased platelet production. It also promotes the proliferation of hepatocytes in response to liver damage. Binding to its receptor formed by IL6ST and either IL11RA1 or IL11RA2, It activates a signaling cascade that promotes cell proliferation. The signaling leads to the activation of intracellular protein kinases and the phosphorylation of STAT3.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com

Web: www.elabscience.com