

# Recombinant Human ROR1 Protein (His Tag)

Catalog Number:PKSH033647



**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

|                                    |  |
|------------------------------------|--|
| <b>Synonyms</b>                    | Inactive tyrosine-protein kinase transmembrane receptor ROR1;Neurotrophic tyrosine kinase;receptor-related 1;ROR1;NTRKR1 |
| <b>Species</b>                     | Human  |
| <b>Expression Host</b>             | HEK293 Cells   |
| <b>Sequence</b>                    | Gln30-Glu403   |
| <b>Accession</b>                   | Q01973   |
| <b>Calculated Molecular Weight</b> | 42.8 kDa   |
| <b>Observed molecular weight</b>   | 60-80 kDa  |
| <b>Tag</b>                         | C-His  |

## Properties

|                       |   |
|-----------------------|---|
| <b>Purity</b>         | > 95 % as determined by reducing SDS-PAGE.  |
| <b>Endotoxin</b>      | < 1.0 EU per µg of the protein as determined by the LAL method.   |
| <b>Storage</b>        | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.             |
| <b>Shipping</b>       | This product is provided as lyophilized powder which is shipped with ice packs.   |
| <b>Formulation</b>    | Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.<br>Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.<br>Please refer to the specific buffer information in the printed manual. |
| <b>Reconstitution</b> | Please refer to the printed manual for detailed information.  |

## Data



> 95 % as determined by reducing SDS-PAGE.

## Background

Receptor tyrosine kinase-like orphan receptor 1 (ROR1); also known as neurotrophic tyrosine kinase; it is a member of the ROR family within receptor tyrosine kinases (RTK) superfamily. Human ROR1 is a type I transmembrane protein with 937 amino acids (aa) in length. It contains a 29 aa signal sequence; a 377 aa extracellular domain (ECD); a 21 aa transmembrane segment; and a 510 aa cytoplasmic region. ROR1 expressed strongly in human heart; lung and kidney; but weakly in the CNS. At developmental stage; it expressed at high levels during early embryonic development. ROR1 has been shown to have very low kinase activity in vitro and is unlikely to function as a tyrosine kinase in vivo. It may act as a receptor for wnt ligand WNT5A which may result in the inhibition of WNT3A-mediated signaling.

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017