A Reliable Research Partner in Life Science and Medicine

Recombinant Human DKK1/Dkk-1 Protein (His Tag)

Catalog No. PKSH033697

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms Dickkopf-related protein 1;Dickkopf-1;Dkk-1;SK

Species Human

Expression Host HEK293 Cells
Sequence Thr32-His266
Accession O94907
Calculated Molecular Weight 27.0 kDa

Calculated Molecular Weight27.0 kDaObserved molecular weight35-50 kDaTagN-His

Bioactivity Not validated for activity

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 0.01 EU per μg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Dickkopf-related protein 1(DKK-1), is a member of the dickkopf family. DKK1 secreted proteins with two cysteine-rich domains separated by a linker region. It antagonizes canonical Wnt signaling by inhibiting LRP5/6 interaction with Wnt

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com

 $Email: \underline{tech support@elabscience.com}$

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

and by forming a ternary complex with the transmembrane protein KREMEN that promotes internalization of LRP5/6. DKKs play an important role in vertebrate development, where they locally inhibit Wnt regulated processes such as anteroposterior axial patterning, limb development, somitogenesis and eye formation. In the adult, Dkks are implicated in bone formation and bone disease, cancer and Alzheimer disease.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com

Web: www.elabscience.com