# Recombinant Human Coagulation Factor X/F10 Protein (Fc

# Tag)

Catalog Number: PKSH033714



Note: Centrifuge before opening to ensure complete recovery of vial contents.

## **Description**

Synonyms Coagulation factor 10;coagulation factor X;FX;FXA

Species Human

**Expression Host** HEK293 Cells **Sequence** Asn32-Lys488

Accession P00742
Calculated Molecular Weight 78.2 kDa

Observed molecular weight 80-120&25 kDa

Tag C-Fc

#### **Properties**

**Purity** > 90 % as determined by reducing SDS-PAGE.

**Endotoxin**  $< 1.0 \text{ EU per } \mu \text{g of the protein as determined by the LAL method.}$ 

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation** Lyophilized from a 0.2 μm filtered solution of 20mM MES, 150mM NaCl, 0.2mM


CaCl<sub>2</sub>, pH 5.5.

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization. Please refer to the specific buffer infor

**Reconstitution** Please refer to the printed manual for detailed information.

#### Data



> 90 % as determined by reducing SDS-PAGE.

## **Background**

F10, also known as Coagulation factor X, belongs to the peptidase S1 family that is synthesized as a 488 amino acid (aa) with a signal peptide and a pro region (residues 1-40). Both the intrinsic and extrinsic pathways activate Factor X to Xa, which consists of light (residues 41-179) and heavy (residues 235-488) chains linked by a disulfide bond. Coagulation factor X is initially synthesized in the liver. The two chains are formed from a single-chain precursor by the excision of two Arg residues and are held together by 1 or more disulfide bonds. Forms a heterodimer with SERPINA5. F10 is a vitamin K-dependent glycoprotein that converts prothrombin to thrombin in the presence of factor Va, calcium and phospholipid during blood clotting.

#### For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: <a href="mailto:www.elabscience.com">www.elabscience.com</a>
Email: <a href="mailto:techsupport@elabscience.com">techsupport@elabscience.com</a>