
Recombinant Human CDO/CDON Protein (His Tag)

Catalog No. PKSH033720

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Synonyms	Cell adhesion molecule-related/down-regulated by oncogenes;CDON;CDO
Species	Human
Expression Host	HEK293 Cells
Sequence	Asp26-Pro943
Accession	Q4KMG0
Calculated Molecular Weight	100.4 kDa
Observed molecular weight	120-135 kDa
Tag	C-His
Bioactivity	Not validated for activity
Properties	
Purity	> 85 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per μ g of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 85 % as determined by reducing SDS-PAGE.

Background

CDO (CAMrelated/down-regulatedby oncogenes) is a member of the Immunoglubulin (Ig) superfamily, Ig/Fibronectin (FN)type III repeat family of cell surface proteins. Human CDO is a type I transmembrane (TM) glycoprotein. It is

For Research Use Only

Toll-free: 1-888-852-8623 Web: <u>www.elabscience.com</u> Tel: 1-832-243-6086 Email: <u>techsupport@elabscience.com</u>

Elabscience®

synthesized as a 1287 amino acid (aa) precursorthat contains a 25 aa signal sequence, a 938 aa extracellular domain (ECD), a 21 aa TM segment and a 303 aa cytoplasmic region. The ECD contains five C2-typeIglikedomains, followed by three FN type III repeats. The ECD of human CDO is 85% aa identical to mouse CDO ECD.CDO is found on muscle precursor and neural progenitor cells of the embryo. It likely promotes muscle differentiation, and contributes to axon guidance andneuronal patterning.

For Research Use Only