

# Recombinant Human BCAM Protein (His Tag)

Catalog Number:PKSH033754



**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

<b>Synonyms</b>	Basal cell adhesion molecule;Aubeger B antigen;B-CAM cell surface glycoprotein;F8/G253 antigen;Lutheran antigen;Lutheran blood group glycoprotein;CD239;BCAM;LU;MSK19
<b>Species</b>	Human
<b>Expression Host</b>	HEK293 Cells
<b>Sequence</b>	Glu32-Ala547
<b>Accession</b>	P50895
<b>Calculated Molecular Weight</b>	57.0 kDa
<b>Observed molecular weight</b>	72-85 kDa
<b>Tag</b>	C-His

## Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

## Data



> 95 % as determined by reducing SDS-PAGE.

## Background

Basal cell adhesion molecule (BCAM, CD239) is an immunoglobulin superfamily protein that arises from alternate splicing of the Lutheran blood group molecule (Lu). The ECD of human BCAM contains two Ig-like V-type domains and three Ig-like C2-type domains. It shares 73% aa sequence identity with the ECDs of mouse and rat BCAM. BCAM is widely expressed in epithelial and endothelial tissues including in the vasculature, kidney glomerulus, small intestine, colon, hair follicle outer root sheath, and basal keratinocytes of the skin during inflammation. BCAM is also expressed on vascular and visceral smooth muscle cells and at the neuromuscular junction of skeletal muscle. BCAM is upregulated on

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

# Recombinant Human BCAM Protein (His Tag)

Catalog Number:PKSH033754



carcinomas, particularly ovarian, sarcomas, astrocytomas, and melanomas. It may mediate intracellular signaling. It cooperates with Integrins  $\beta 1$  and  $\alpha V\beta 3$  as an adhesion receptor for Laminins which contain the  $\alpha 5$  chain. The Lutheran isoform is aberrantly phosphorylated in erythroid disorders and can enhance Lamininmediated adhesion of erythrocytes to vascular endothelial cells.

---

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Tel: 1-832-243-6086

Fax: 1-832-243-6017

Web: [www.elabscience.com](http://www.elabscience.com)

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)