

Recombinant Human IL3RA/CD123 Protein (Fc & Avi Tag)

Catalog No. PKSH033800

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Interleukin-3 receptor subunit alpha;IL-3 receptor subunit alpha;IL-3R subunitalpha;IL-3R-alpha;IL-3RA
Species	Human
Expression Host	HEK293 Cells
Sequence	Thr19-Arg305
Accession	P26951
Calculated Molecular Weight	61.8 kDa
Observed molecular weight	80-100 kDa
Tag	C-Fc-Avi
Bioactivity	Not validated for activity

Properties

Purity	> 90 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 90 % as determined by reducing SDS-PAGE.

Background

CD123; also known as Interleukin-3 receptor subunit alpha; belongs to the type I cytokine receptor family. In mouse;

For Research Use Only

there are two classes of high-affinity IL3 receptors. One contains an IL3-specific beta subunit and the other contains the beta subunit also shared by high-affinity IL5 and GM-CSF receptors. CD123 stimulates the proliferation and differentiation of hemopoietic cells including the pluripotent hematopoietic stem cells as well as various lineage-committed cells. CD123 is a heterodimer consisting of an alpha and a beta subunit. The alpha subunit alone binds IL-3 with low affinity. The beta subunit does not bind IL-3 by itself but is required for the high-affinity binding of IL-3 to the heterodimeric receptor complex.