

## Recombinant Human/Mouse/Rat BDNF Protein

**Catalog No.** PKSH033808

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Synonyms</b>	Brain-Derived Neurotrophic Factor;BDNF;Abrineurin
<b>Species</b>	Human/Mouse/Rat
<b>Expression Host</b>	E.coli
<b>Sequence</b>	His129-Arg247
<b>Accession</b>	P23560
<b>Calculated Molecular Weight</b>	14.5 kDa
<b>Observed molecular weight</b>	14 kDa
<b>Tag</b>	C-His
<b>Bioactivity</b>	Measure by its ability to induce proliferation in BaF3 cells transfected with TrkB. The ED <sub>50</sub> for this effect is < 2 ng/mL.

### Properties

<b>Purity</b>	> 98 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 0.1 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile 20 mM sodium citrate, 0.2 M NaCl, pH 3.5. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 98 % as determined by reducing SDS-PAGE.

### Background

Brain-Derived Neurotrophic Factor (BDNF) is a member of the neurotrophin family. Along with other structurally related

### For Research Use Only

neurotrophic factors NGF; NT-3 and NT-4; BDNF binds with high affinity to the TrkB kinase receptor. It also binds with the LNGFR (for low-affinity nerve growth factor receptor; also known as p75). BDNF promotes the survival; growth and differentiation of neurons. It serves as a major regulator of synaptic transmission and plasticity at adult synapses in many regions of the CNS. BDNF expression is altered in neurodegenerative disorders such as Parkinson's and Alzheimer's disease.