A Reliable Research Partner in Life Science and Medicine

Recombinant Mouse VCAM1 Protein (Fc Tag)

Catalog No. PKSM041260

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms CD106;Vascular cell adhesion protein 1;Vcam1;L1CAM;VCMA1;Vcam-1

Species Mouse

Expression Host HEK293 Cells **Sequence** Phe25-Glu698

AccessionP29533Calculated Molecular Weight101.4 kDaObserved molecular weight100-130 kDa

Tag C-Fc

Bioactivity Not validated for activity

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per ug of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Vascular cell adhesion molecule 1 (VCAM-1) is a cell surface protein belonging to the immunoglobulin superfamily, the protein is expressed by activated endothelial cells and certain leukocytes (such as macrophages). IL-1 beta, IL-4, TNF-

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com

 $Email: \underline{tech support@elabscience.com}$

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

alpha and IFN-gamma induced the expression of VCAM-1. The human and mouse VCAM-1 proteins share approximately 76% amino acid similarity. Mouse VCAM-1 is Important in cell-cell recognition. it appears to function in leukocyte-endothelial cell adhesion, and interacts with integrin alpha-4/beta-1 (ITGA4/ITGB1) on leukocytes, and mediates both adhesion and signal transduction.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com

Web: www.elabscience.com