
# Recombinant Mouse TRAIL R2/TNFRSF10B Protein (Fc Tag)



Catalog Number: PKSM041271

Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description                 |  |
|-----------------------------|--|
| Synonyms                    | Tumor Necrosis Factor Receptor Superfamily Member 10B;Death Receptor 5;TNF-<br>Related Apoptosis-Inducing Ligand Receptor 2;TRAIL Receptor 2;TRAIL-<br>R2;CD262;TNFRSF10B;DR5;KILLER;TRAILR2;TRICK2;ZTNFR9   |
| Species                     | Mouse  |
| Expression Host             | HEK293 Cells   |
| Sequence                    | Asn53-Ser177   |
| Accession                   | Q9QZM4   |
| Calculated Molecular Weight | 40.9 kDa   |
| Observed molecular weight   | 50-75 kDa  |
| Tag                         | C-Fc   |
| Properties                  |  |
| Purity                      | > 95 % as determined by reducing SDS-PAGE.   |
| Endotoxin                   | < 1.0 EU per µg of the protein as determined by the LAL method.  |
| Storage                     | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.                            |
| Shipping                    | This product is provided as lyophilized powder which is shipped with ice packs.  |
| Formulation                 | Lyophilized from a 0.2 µm filtered solution of 20mM PB,150mM NaCl,pH7.4.<br>Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as<br>protectants before lyophilization.<br>Please refer to the specific buffer information in the printed manual |
| Reconstitution              | Please refer to the printed manual for detailed information.   |
| Data                        |  |



> 95 % as determined by reducing SDS-PAGE.

### Background

Mouse tumor necrosis factor receptor superfamily member 10B (TNFRSF10B) is a member of the TNFR family which contains 1 death domain and 3 TNFR-Cys repeats. TNFRSF10B exhibits high structural and functional homology to TRAIL-R1 (DR-4). TNFRSF10B is highly expressed in heart, lung, lymphocytes, spleen and kidney. In addition, it is regulated by the tumor suppressor p53. TNFRSF10B is the receptor for the cytotoxic ligand TNFSF10/TRAIL. It promotes the activation of NF-kappa-B and is essential for ER stress-induced apoptosis. The adapter molecule FADD recruits caspase-8 to the activated receptor. The resulting death-inducing signaling complex (DISC) performs caspase-8

### For Research Use Only

A Reliable Research Partner in Life Science and Medicine Toll-free: 1-888-852-8623 Tel: 1-832-243-6086

## Recombinant Mouse TRAIL R2/TNFRSF10B Protein (Fc Tag)



Catalog Number:PKSM041271

proteolytic activation which initiates the subsequent cascade of caspases mediating apoptosis.

#### For Research Use Only

A Reliable Research Partner in Life Science and Medicine Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web: www.elabscience.com Email: techsupport@elabscience.com