

Recombinant Mouse PDGF-BB Protein (His Tag)

Catalog No. PKSM041293

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Platelet-Derived Growth Factor Subunit B;PDGF Subunit B;PDGF-2;Platelet-Derived Growth Factor B Chain;Platelet-Derived Growth Factor Beta Polypeptide;Proto-Oncogene c-Sis;Becaplermin;PDGFB;PDGF2;SIS
Species	Mouse
Expression Host	E.coli
Sequence	Ser82-Thr190
Accession	AAH53430.1
Calculated Molecular Weight	13.4 kDa
Observed molecular weight	15 kDa
Tag	C-His
Bioactivity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of 4mM HCl. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Platelet-Derived Growth Factor Subunit B (PDGFB) belongs to the PDGF/VEGF growth factor family. Platelet-derived growth factor is a potent mitogen for cells of mesenchymal origin. PDGFB can exist either as a homodimer (PDGF-BB) or as a heterodimer with the platelet-derived growth factor alpha polypeptide (PDGF-AB), where the dimers are connected by disulfide bonds. As growth factor, it plays an essential role in the regulation of embryonic development, cell proliferation, cell migration, survival and chemotaxis. It is required for normal proliferation and recruitment of pericytes and vascular smooth muscle cells in the central nervous system, skin, lung, heart and placenta. PDGFB also plays an important role in wound healing.