

Recombinant Mouse IGFBP-7/IGFBP7 Protein (His Tag)

Catalog No. PKSM041324

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Insulin-like growth factor-binding protein 7;IGFBP7;IGF-binding protein 7;IGFBP-rP1;MAC25 protein;Tumor-derived adhesion factor;TAF;Igfbp7;IBP-7;AGM;Fstl2;Mac25
Species	Mouse
Expression Host	HEK293 Cells
Sequence	Ser26-Leu281
Accession	Q61581
Calculated Molecular Weight	27.2 kDa
Observed molecular weight	32-35 kDa
Tag	C-His
Bioactivity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, 150mM NaCl, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Insulin-like growth factor-binding protein 7(IGFBP-7) is a secreted glycosylated protein that contains three protein domain modules. IGFBP7 contains an N-terminal IGFBP domain, followed by a Kazal-type serine proteinase inhibitor domain and a C-terminal immunoglobulin-like C2-type domain. Human and mouse IGFBP7 are highly homologous and share 94% aa sequence identity. It is expressed in many normal tissues and in cancer cells. It is abundantly expressed in high endothelial venules (HEVs) of blood vessels in the secondary lymphoid tissues. It binds IGF and insulin with very low affinity and has been shown to enhance the mitogenic actions of IGF and insulin. IGFBP7 also has IGF/insulin-independent activities. It interacts with heparan sulfate proteoglycans, type IV collagen, and specific chemokines. It supports weak cell adhesion, promotes cell spreading on type IV collagen, and stimulates the production of the potent vasodilator PGI₂. It modulates tumor cell growth and has also been implicated in angiogenesis.