## **Recombinant Mouse ICOS Ligand/ICOSL Protein (His**

### Tag)

#### Catalog Number: PKSM041331



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Descri	4.0
LIACOPT	ntion
	747 (17)

Synonyms B7 homolog 2;B7-H2;B7-like protein Gl50;B7RP-1LICOS;CD275;CD275

antigen;ICOS ligand;ICOSL;ICOS-L;inducible T-cell co-stimulator

ligand;B7RP-1;CD275;GL50;ICOSL;ICOSLG;B7H2;B7RP1;ICOS Ligand

**Species** Mouse

Expression Host
Sequence
Glu47-Lys279
Accession
Q9JHJ8
Calculated Molecular Weight
Observed molecular weight
Tag
HEK293 Cells
Glu47-Lys279
49JHJ8
27.1 kDa
40-60 kDa
C-His

#### **Properties**

**Purity** > 95 % as determined by reducing SDS-PAGE.

**Endotoxin** < 1.0 EU per µg of the protein as determined by the LAL method.

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation** Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

#### Data



> 95 % as determined by reducing SDS-PAGE.

#### **Background**

Mouse ICOS ligand(B7-H2) is an approximately transmembrane glycoprotein in the B7 family of immune regulatory molecules. B7-H2 is expressed on antigen presenting cells such as B cells, macrophages, monocytes, and dendritic cells. It binds to ICOS on activated T cells, leading to both positive and negative effects on immune responses including its own down-regulation. The B7-H2 interaction with ICOS is costimulatory for T cell proliferation as well as the development of B cells, plasma cells, follicular helper T cells and germinal centers. B7-H2 contributes to the development of allergic asthma by enhancing Th2 biased immune responses, limiting Th17 responses, and promoting eosinophilic infiltration into

#### For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com

# **Recombinant Mouse ICOS Ligand/ICOSL Protein (His Tag)**



Catalog Number:PKSM041331

the lung. Its activation of ICOS on Treg limits pulmonary inflammation and airway hyperresponsiveness, promotes the development of inhalational tolerance, and impairs antitumor immunity. In the thyroid, B7-H2 is up-regulated on thyrocytes during inflammation and promotes their proliferation and production of thryoid hormones.

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017
Web: www.elabscience.com Email: techsupport@elabscience.com