

Recombinant Mouse NTPDase 2/ENTPD2 Protein (His Tag)

Catalog No. PKSM041340

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Ectonucleoside triphosphate diphosphohydrolase 2;Entpd2;Ecto-Nucleoside Triphosphate Diphosphohydrolase 2
Species	Mouse
Expression Host	HEK293 Cells
Sequence	Cys26-Ser462
Accession	O55026
Calculated Molecular Weight	49.2 kDa
Observed molecular weight	63-90 kDa
Tag	C-His
Bioactivity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.
Shipping	This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel packs. Upon receipt, store it immediately at < -20°C.
Formulation	Supplied as a 0.2 µm filtered solution of 50mM Tris-HCl, 10mM CaCl ₂ , 150mM NaCl, 10% Glycerol, pH 7.5.
Reconstitution	Not Applicable

Data



> 95 % as determined by reducing SDS-PAGE.

Background

CD39L1 protein (ENTPD2 or NTPDase2) is a member of the ecto-nucleoside triphosphate diphosphohydrolase family which the main role is termination of purinergic signaling. CD39L1 gene encodes a precursor protein with 495 amino acid residues which generates a 437 amino acid residues mature protein after processing. It is an ecto-nucleotidase that found on the surface of vascular adventitial cells and accessory vascular cells. CD39L1 is a Ca²⁺- and Mg²⁺-dependent enzyme

For Research Use Only

that activates platelets by preferentially converting ATP to ADP. CD39L1 plays a role in regulating thrombosis and inflammation which is considered to be a therapeutic target for thromboregulation and the treatment of vascular inflammation. Alternative splicing of CD39L1 gene results in multiple transcript variants.