

# Recombinant Mouse CD36/SCARB3 Protein (AVI & His Tag)



Catalog Number: PKSM041371

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

<b>Synonyms</b>	Glycoprotein IIb; GPIIIB; PAS IV; PAS-4; Platelet glycoprotein IV; GPIV; CD36
<b>Species</b>	Mouse
<b>Expression Host</b>	HEK293 Cells
<b>Sequence</b>	Gly30-Lys439
<b>Accession</b>	Q08857
<b>Calculated Molecular Weight</b>	49.0 kDa
<b>Observed molecular weight</b>	85-120 kDa
<b>Tag</b>	C-Avi-His

## Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01 % Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

## Data



> 95 % as determined by reducing SDS-PAGE.

## Background

Platelet Glycoprotein 4(CD36) is belongs to the class B scavenger receptor family. The molecule CD36 is synthesized as a 472 amino acid (aa) protein that contains a 6 aa N-terminal cytoplasmic domain, a 22 aa N-terminal transmembrane segment, a 420 aa extracellular “loop”, a 22 aa C-terminal transmembrane segment, and a 9 aa C-terminal cytoplasmic tail. Both cytoplasmic tails are palmitoylated, with the C-terminal tail involved in oxidized LDL binding. With respect to the extracellular loop, the N-terminal region is believed to bind both thrombospondin-1 and Plasmodium-infected erythrocytes. Other ligands for CD36 include long-chain fatty acids, collagen, phospholipids and apoptotic cells. Cells known to express CD36 include capillary endothelium, adipocytes, skeletal muscle cells, intestinal epithelium, smooth muscle cells and hematopoietic cells such as RBC's, platelets and monocytes. On the surface of cells, CD36 is suggested to

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

# Recombinant Mouse CD36/SCARB3 Protein (AVI & His Tag)



Catalog Number: PKSM041371

---

exist as a dimer in response to ligation (7). CD36 is reported to regulate fatty uptake, act as an angiogenic with TSP-1, and participate in the clearance of apoptotic phagocytes.

---

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017