

# Recombinant Rat CD73/NT5E Protein (His Tag)

Catalog Number:PKSR030462



**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

<b>Synonyms</b>	CD73;E5NT;Nucleotidase;NT5E;ecto-5'-nucleotidase;5'-Nucleotidase
<b>Species</b>	Rat
<b>Expression Host</b>	HEK293 Cells
<b>Sequence</b>	Trp29-Lys549
<b>Accession</b>	P21588
<b>Calculated Molecular Weight</b>	59.0 kDa
<b>Observed molecular weight</b>	70-90 kDa
<b>Tag</b>	C-His

## Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.
<b>Shipping</b>	This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel packs. Upon receipt, store it immediately at < - 20°C.
<b>Formulation</b>	Supplied as a 0.2 µm filtered solution of 20mM Tris-HCl, 120mM NaCl, 4mM CaCl <sub>2</sub> , 20% Glycerol, pH 7.5.
<b>Reconstitution</b>	Not Applicable

## Data



> 95 % as determined by reducing SDS-PAGE.

## Background

CD73, also known as Ecto-5'-nucleotidase, is an ectoenzyme that is attached to the cell membrane by a glycosyl phosphatidylinositol anchor. The enzyme is expressed by most cell types. The 5' -Nucleotidase activity of CD73 converts extracellular nucleoside-5' -monophosphates to nucleosides. CD73 is one of several enzymes responsible for the production of extracellular adenosine, a signaling molecule that is involved in responses to inflammation and tissue injury. CD73 hydrolyzes extracellular nucleotides into membrane permeable nucleosides.

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017