

FUT3 Polyclonal Antibody

Catalog No. E-AB-40309

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Human
Immunogen	Recombinant Human Galactoside 3(4)-L-fucosyltransferase protein
Host	Rabbit
Isotype	IgG
Purification	Antigen Affinity Purification
Conjugation	Unconjugated
Buffer	PBS with 0.05% Proclin300 and 50% glycerol, pH7.4.


Applications Recommended Dilution

IHC 1:100-1:300


Data




Immunohistochemistry of paraffin-embedded Human kidney using FUT3 Polyclonal Antibody at dilution of 1:200.



Immunohistochemistry of paraffin-embedded Human stomach cancer using FUT3 Polyclonal Antibody at dilution of 1:200.



Immunohistochemistry of paraffin-embedded Human Colorectal cancer using FUT3 Polyclonal Antibody at dilution of 1:200(100×).



Immunohistochemistry of paraffin-embedded Human Colorectal cancer using FUT3 Polyclonal Antibody at dilution of 1:200(400×).

Preparation & Storage

Storage Store at -20°C. Avoid freeze / thaw cycles.

Background

The Lewis histo-blood group system comprises a set of fucosylated glycosphingolipids that are synthesized by exocrine

For Research Use Only

epithelial cells and circulate in body fluids. The glycosphingolipids function in embryogenesis, tissue differentiation, tumor metastasis, inflammation, and bacterial adhesion. They are secondarily absorbed to red blood cells giving rise to their Lewis phenotype. This gene is a member of the fucosyltransferase family, which catalyzes the addition of fucose to precursor polysaccharides in the last step of Lewis antigen biosynthesis. It encodes an enzyme with alpha(1,3)-fucosyltransferase and alpha(1,4)-fucosyltransferase activities. Mutations in this gene are responsible for the majority of Lewis antigen-negative phenotypes. Multiple alternatively spliced variants, encoding the same protein, have been found for this gene.