

Histone H3 Monoclonal Antibody

Catalog No. E-AB-22003

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Human, Mouse, Rat
Immunogen	Recombinant Protein
Host	Mouse
Isotype	IgG
Clone	5B1
Purification	Protein A purification
Conjugation	Unconjugated
Buffer	PBS with 0.02% sodium azide and 50% glycerol pH 7.4.

Applications Recommended Dilution

WB	1:500-1:2000
IF	1:100-500


Data



www.elabscience.com

Western Blot analysis of 1) HeLa, 2) Raw264.7, 3) Mouse brain, 4) Rat brain using Histone H3 Monoclonal Antibody at dilution of 1:5000.

Observed Mw:15kDa
Calculated Mw:15kDa



www.elabscience.com

Immunofluorescence analysis of Rat liver tissue using Histone H3 Monoclonal Antibody at dilution of 1:200.

Preparation & Storage

Storage Store at -20°C. Avoid freeze / thaw cycles.

Background

Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.

For Research Use Only