

CDKL4 Polyclonal Antibody

Catalog No. E-AB-17783

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Human, Mouse
Immunogen	Synthetic peptide of human CDKL4
Host	Rabbit
Isotype	IgG
Purification	Antigen affinity purification
Conjugation	Unconjugated
Buffer	PBS with 0.05% NaN ₃ and 40% Glycerol, pH7.4

Applications Recommended Dilution


WB	1:500-1:2000
IHC	1:30-1:150

Data




Western blot analysis of Human heart tissue lysate using CDKL4 Polyclonal Antibody at dilution of 1:550

Observed Mw: Refer to figures
Calculated Mw: 43 kDa



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using CDKL4 Polyclonal Antibody at dilution of 1:45 (×200)



Immunohistochemistry of paraffin-embedded Human ovarian cancer tissue using CDKL4 Polyclonal Antibody at dilution of 1:45 (×200)

Preparation & Storage

For Research Use Only

Storage Store at -20°C. Avoid freeze / thaw cycles.

Background

Cell cycle progression is controlled in part by a family of cyclin proteins and cyclin dependent kinases (Cdks). Cdk proteins work in concert with the cyclins to phosphorylate key substrates involved in each phase of cell cycle progression. Another family of proteins, Cdk inhibitors, also plays a role in regulating the cell cycle by binding to cyclin-Cdk complexes and modulating their activity. Cdks are considered potential targets for anti-cancer therapy due to their involvement with cell cycle regulation. Cdks are also involved in the regulation of transcription and mRNA processing. CdkL4 (cyclin-dependent kinase-like 4) is a 315 amino acid protein that belongs to the CMGC Ser/Thr protein kinase family and may be involved in cell cycle regulation.