
## **GRIA2** Polyclonal Antibody

Catalog Number: E-AB-18121



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human, Mouse, Rat
Immunogen	Synthetic peptide of human GRIA2
Host	Rabbit
Isotype	IgG
Purification	Antigen affinity purification
Conjugation	Unconjugated
Formulation	PBS with 0.05% NaN3 and 40% Glycerol,pH7.4
Applications	Recommended Dilution
IHC	1:50-1:300
ELISA	1:5000-1:10000
Data	



Immunohistochemistry of paraffin-embedded Human tonsil tissue using GRIA2 Polyclonal Antibody at dilution of 1:50(×200)

## **Preparation & Storage**

Storage

Store at -20°C. Avoid freeze / thaw cycles.

## Background

Glutamate receptors are the predominant excitatory neurotransmitter receptors in the mammalian brain and are activated in a variety of normal neurophysiologic processes. This gene product belongs to a family of glutamate receptors that are sensitive to alpha-amino-3-hydroxy-5-methyl-4-isoxazole propionate (AMPA), and function as ligand-activated cation channels. These channels are assembled from 4 related subunits, GRIA1-4. The subunit encoded by this gene (GRIA2) is subject to RNA editing (CAG->CGG; Q->R) within the second transmembrane domain, which is thought to render the channel impermeable to Ca(2+). Human and animal studies suggest that pre-mRNA editing is essential for brain function, and defective GRIA2 RNA editing at the Q/R site may be relevant to amyotrophic lateral sclerosis (ALS) etiology. Alternative splicing, resulting in transcript variants encoding different isoforms, (including the flip and flop isoforms that vary in their signal transduction properties), has been noted for this gene.

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web: www.elabscience.com Email: techsupport@elabscience.com