

HDAC1 Polyclonal Antibody

Catalog No. E-AB-18230

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Human, Mouse, Rat
Immunogen	Fusion protein of human HDAC1
Host	Rabbit
Isotype	IgG
Purification	Antigen affinity purification
Conjugation	Unconjugated
Buffer	PBS with 0.05% NaN ₃ and 40% Glycerol,pH7.4


Applications Recommended Dilution

WB	1:1000-1:5000
IHC	1:50-1:200

Data



Western blot analysis of 293T 231 Jurkat and RAW264.7 cell lysates using HDAC1 Polyclonal Antibody at dilution of 1:1000
Observed Mw:Refer to figures
Calculated Mw:55 kDa



Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using HDAC1 Polyclonal Antibody at dilution of 1:45(×200)

Preparation & Storage

Storage Store at -20°C. Avoid freeze / thaw cycles.

Background

Histone acetylation and deacetylation, catalyzed by multisubunit complexes, play a key role in the regulation of eukaryotic gene expression. The protein encoded by this gene belongs to the histone deacetylase/acuc/apha family and is a component of the histone deacetylase complex. It also interacts with retinoblastoma tumor-suppressor protein and this complex is a key element in the control of cell proliferation and differentiation. Together with metastasis-associated protein-2, it deacetylates p53 and modulates its effect on cell growth and apoptosis.

For Research Use Only