## PHF21A Polyclonal Antibody

Catalog Number: E-AB-18538



*Note:* Centrifuge before opening to ensure complete recovery of vial contents.

## Description

Reactivity Human, Mouse

**Immunogen** Fusion protein of human PHF21A

Host Rabbit
Isotype IgG

**Purification** Antigen affinity purification


Conjugation Unconjugated

**Formulation** PBS with 0.05% NaN3 and 40% Glycerol,pH7.4

## **Applications** Recommended Dilution


WB 1:500-1:2000 IHC 1:40-1:200 ELISA 1:5000-1:10000

#### Data




Western blot analysis of 293T Hela and A375 cell lysates using PHF21A Polyclonal Antibody at dilution of 1:400

Observed Mw:Refer to figures Calculated Mw:75 kDa



Immunohistochemistry of paraffin-embedded Human prost at e cancer tissue using PHF21A Polyclonal Antibody at dilution of 1:50(×200)



Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using PHF21A Polyclonal Antibody at dilution of 1:50(×200)

## **Preparation & Storage**

**Storage** Store at -20°C. Avoid freeze / thaw cycles.

### **Background**

#### For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: <u>www.elabscience.com</u> Email: <u>techsupport@elabscience.com</u>

# PHF21A Polyclonal Antibody

Catalog Number: E-AB-18538



Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. PHF21A (PHD finger protein 21A), also known as BRAF35-HDAC complex protein BHC80, is a 680 amino acid nuclear protein that contains one PHD-type zinc finger and one A.T hook DNA-binding domain, suggesting involvement in transcriptional regulation events. PHF21A is a component of the BHC complex, which is responsible for repressing transcription of neuron-specific genes in non-neuronal cells. The BHC complex acts as a chromatin modifier that deacetylates and demethylates specific sites on histones. PHF21A may act as a scaffold within the BHC complex. Predominantly expressed in brain, three isoforms of PHF21A exist as a result of alternative splicing events.

#### For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com