

CA7 Polyclonal Antibody

Catalog No. E-AB-18577

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Human, Mouse
Immunogen	Full length fusion protein
Host	Rabbit
Isotype	IgG
Purification	Antigen affinity purification
Conjugation	Unconjugated
Buffer	PBS with 0.05% NaN ₃ and 40% Glycerol, pH7.4


Applications Recommended Dilution

WB	1:500-1:2000
IHC	1:50-1:300


Data



Western blot analysis of Human urinary bladder tissue Mouse heart tissue Mouse kidney tissue and RAW264.7 cell lysates using CA7 Polyclonal Antibody at dilution of 1:500
Observed Mw:Refer to figures
Calculated Mw:30 kDa



Immunohistochemistry of paraffin-embedded Human gastric cancer tissue using CA7 Polyclonal Antibody at dilution of 1:50(×200)



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using CA7 Polyclonal Antibody at dilution of 1:50(×200)

Preparation & Storage

For Research Use Only

Storage Store at -20°C. Avoid freeze / thaw cycles.

Background

Carbonic anhydrases are a large family of zinc metalloenzymes that catalyze the reversible hydration of carbon dioxide. They participate in a variety of biological processes, including respiration, calcification, acid-base balance, bone resorption, and the formation of aqueous humor, cerebrospinal fluid, saliva, and gastric acid. They show extensive diversity in tissue distribution and in their subcellular localization. The cytosolic protein encoded by this gene is predominantly expressed in the salivary glands. Alternative splicing in the coding region results in multiple transcript variants encoding different isoforms.

For Research Use Only