

## CCT8L2 Polyclonal Antibody

**Catalog No.** E-AB-18611

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

|                     |                                                         |
|---------------------|---------------------------------------------------------|
| <b>Reactivity</b>   | Human                                                   |
| <b>Immunogen</b>    | Fusion protein of human CCT8L2                          |
| <b>Host</b>         | Rabbit                                                  |
| <b>Isotype</b>      | IgG                                                     |
| <b>Purification</b> | Antigen affinity purification                           |
| <b>Conjugation</b>  | Unconjugated                                            |
| <b>Buffer</b>       | PBS with 0.05% NaN <sub>3</sub> and 40% Glycerol, pH7.4 |


### Applications Recommended Dilution

|            |            |
|------------|------------|
| <b>IHC</b> | 1:30-1:150 |
|------------|------------|

### Data



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using CCT8L2 Polyclonal Antibody at dilution of 1:45(×200)



Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using CCT8L2 Polyclonal Antibody at dilution of 1:45(×200)

### Preparation & Storage

**Storage** Store at -20°C. Avoid freeze / thaw cycles.

### Background

Possible molecular chaperone; assists the folding of proteins upon ATP hydrolysis.

### For Research Use Only