
BAG2 Polyclonal Antibody

Catalog Number: E-AB-19397



Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description	
Reactivity	Human
Immunogen	Synthetic peptide of human BAG2
Host	Rabbit
Isotype	IgG
Purification	Antigen affinity purification
Conjugation	Unconjugated
Formulation	PBS with 0.05% NaN3 and 40% Glycerol,pH7.4
Applications	Recommended Dilution
WB	1:500-1:2000
IHC	1:30-1:150
ELISA	1:5000-1:10000
Data	



Western blot analysis of Human fetal muscle tissue HEPG2 Jurkat Hela A431 and A549 cell using BAG2 Polyclonal Antibody at dilution of 1:550 **Observed Mw:Refer to figures Calculated Mw:24 kDa**



Immunohistochemistry of paraffin-embedded Human tonsil tissue using BAG2 Polyclonal Antibody at dilution of 1:40(×200)



Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using BAG2 Polyclonal Antibody at dilution of 1:40(×200)

Preparation & Storage

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Background

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Web: <u>www.elabscience.com</u> Tel: 1-832-243-6086 Email: <u>techsupport@elabscience.com</u>

BAG2 Polyclonal Antibody

Catalog Number: E-AB-19397



BAG proteins compete with Hip for binding to the Hsc70/Hsp70 ATPase domain and promote substrate release. All the BAG proteins have an approximately 45-amino acid BAG domain near the C terminus but differ markedly in their N-terminal regions. The predicted BAG2 protein contains 211 amino acids. The BAG domains of BAG1, BAG2, and BAG3 interact specifically with the Hsc70 ATPase domain in vitro and in mammalian cells. All 3 proteins bind with high affinity to the ATPase domain of Hsc70 and inhibit its chaperone activity in a Hip-repressible manner.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine
Toll-free: 1-888-852-8623 Tel: 1-832-243-6086
Web: www.elabscience.com Email: techsupport@elabscience.com