

SPTA1 Polyclonal Antibody

Catalog No. E-AB-19527

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Human, Mouse
Immunogen	Synthetic peptide of human SPTA1
Host	Rabbit
Isotype	IgG
Purification	Antigen affinity purification
Conjugation	Unconjugated
Buffer	PBS with 0.05% NaN ₃ and 40% Glycerol, pH7.4

Applications

Recommended Dilution


WB	1:500-1:2000
IHC	1:60-1:450

Data



Western blot analysis of Mouse heart tissue lysate using SPTA1 Polyclonal Antibody at dilution of 1:550

Observed Mw: Refer to figures
Calculated Mw: 280 kDa



Immunohistochemistry of paraffin-embedded Human gastric cancer tissue using SPTA1 Polyclonal Antibody at dilution of 1:80 (x200)

Preparation & Storage

Storage Store at -20°C. Avoid freeze / thaw cycles.

Background

Spectrin is an actin crosslinking and molecular scaffold protein that links the plasma membrane to the actin cytoskeleton, and functions in the determination of cell shape, arrangement of transmembrane proteins, and organization of organelles. It is a tetramer made up of alpha-beta dimers linked in a head-to-head arrangement. This gene is one member of a family of alpha-spectrin genes. The encoded protein is primarily composed of 22 spectrin repeats which are involved in dimer formation. It forms weaker tetramer interactions than non-erythrocytic alpha spectrin, which may increase the plasma membrane elasticity and deformability of red blood cells. Mutations in this gene result in a variety of hereditary red blood cell disorders, including elliptocytosis type 2, pyropoikilocytosis, and spherocytic hemolytic anemia.

For Research Use Only