# Phospho-PI 3 kinase p85 alpha /gamma (Tyr467/199) **Polyclonal Antibody**



Catalog Number: E-AB-20966 2 Publications

Note: Centrifuge before opening to ensure complete recovery of vial contents.

### **Description**

Reactivity Human, Mouse, Rat, Monkey

Synthesized peptide derived from human PI 3-kinase p85/p55 around the **Immunogen** 

phosphorylation site of Tyr467/199

Host Rabbit IgG **Isotype** 


**Purification** Affinity purification Conjugation **Unconjugated** 

PBS with 0.02% sodium azide, 0.5% protective protein and 50% glycerol, pH7.4 **Formulation** 

#### **Applications Recommended Dilution**

WB 1:500-1:2000 **IHC** 1:100-1:300 IF 1:200-1:1000 **ELISA** 1:10000

### Data



Western Blot analysis of various cells using Phospho-PI 3 kinase p85 alpha /gamma (Tyr467/199) Polyclonal Antibody at dilution of 1:1000

> Observed Mw:55+85kDa Calculated Mw:54+83kDa



Immunohistochemistry of paraffin-embedded Human colon tissue using Phospho-PI 3 kinase p85 alpha /gamma (Tyr467/199) Polyclonal Antibody at dilution of 1:200



Immunofluorescence analysis of Rat spleen tissue using Phospho-PI 3 kinase p85 alpha /gamma (Tyr467/199) Polyclonal Antibody at dilution of 1:200

## **Preparation & Storage**

### For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com

# Phospho-PI 3 kinase p85 alpha /gamma (Tyr467/199) Polyclonal Antibody



Catalog Number: E-AB-20966 2 Publications

Storage Store at -20°C. Avoid freeze / thaw cycles.

### **Background**

The enzyme phosphatidylinositol 3 kinase (PI3 kinase) is a lipid kinase that generates phosphatidylinositol 3, 4, 5-triphosphate in response to receptor activation in many signal transduction pathways. Class IA PI3Ks exist as a heterodimer of a catalytic 110 kDa (p110) and a regulatory p85 subunit (e.g. p85 alpha). p85 alpha is an adaptor molecule that regulates the activity of the catalytic p110 subunit by binding to phosphorylated receptor tyrosine kinases (RTKs) through its SH2 domain and mediating the interaction between p110 and the plasma membrane. p85 alpha is necessary for insulin-stimulated increase in glucose uptake and glycogen synthesis in insulin-sensitive tissues.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: <u>www.elabscience.com</u> Email: <u>techsupport@elabscience.com</u>