
## **CD8A** Polyclonal Antibody

Catalog Number: E-AB-19675



*Note:* Centrifuge before opening to ensure complete recovery of vial contents.

| Description  |  |
|--------------|--|
| Reactivity   | Human                                      |
| Immunogen    | Synthetic peptide of human CD8A            |
| Host         | Rabbit                                     |
| Isotype      | IgG  |
| Purification | Antigen affinity purification              |
| Conjugation  | Unconjugated                               |
| Formulation  | PBS with 0.05% NaN3 and 40% Glycerol,pH7.4 |
| Applications | Recommended Dilution                       |
| IHC          | 1:50-1:300                                 |
| ELISA        | 1:5000-1:10000                             |
| Data         |  |



Immunohistochemistry of paraffin-embedded Human prost at e cancer tissue using CD8A Polyclonal Antibody at dilution of 1:100(×200)

## **Preparation & Storage**

Storage

Store at -20°C. Avoid freeze / thaw cycles.

## Background

The CD8 antigen is a cell surface glycoprotein found on most cytotoxic T lymphocytes that mediates efficient cell-cell interactions within the immune system. The CD8 antigen acts as a coreceptor with the T-cell receptor on the T lymphocyte to recognize antigens displayed by an antigen presenting cell in the context of class I MHC molecules. The coreceptor functions as either a homodimer composed of two alpha chains or as a heterodimer composed of one alpha and one beta chain. Both alpha and beta chains share significant homology to immunoglobulin variable light chains. This gene encodes the CD8 alpha chain. Multiple transcript variants encoding different isoforms have been found for this gene.

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web: www.elabscience.com Email: techsupport@elabscience.com