

IDUA Polyclonal Antibody

Catalog No. E-AB-52010

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Human, Mouse
Immunogen	Synthetic peptide of human IDUA
Host	Rabbit
Isotype	IgG
Purification	Antigen affinity purification
Conjugation	Unconjugated
Buffer	PBS with 0.05% NaN ₃ and 40% Glycerol, pH7.4

Applications


Recommended Dilution

IHC 1:30-1:150

Data



Immunohistochemistry of paraffin-embedded Human tonsil tissue using IDUA Polyclonal Antibody at dilution of 1:30(×200)



Immunohistochemistry of paraffin-embedded Human colorectal cancer tissue using IDUA Polyclonal Antibody at dilution of 1:30(×200)

Preparation & Storage

Storage Store at -20°C. Avoid freeze / thaw cycles.

Background

This gene encodes an enzyme that hydrolyzes the terminal alpha-L-iduronic acid residues of two glycosaminoglycans, dermatan sulfate and heparan sulfate. This hydrolysis is required for the lysosomal degradation of these glycosaminoglycans. Mutations in this gene that result in enzymatic deficiency lead to the autosomal recessive disease mucopolysaccharidosis type I (MPS I).

For Research Use Only