Phospho-MAP3K7 (Thr187) Polyclonal Antibody

Catalog Number: E-AB-21065 1 Publications



Note: Centrifuge before opening to ensure complete recovery of vial contents.

		- 40	
Desc	\mathbf{r}	$\mathbf{n}\mathbf{T}\mathbf{I}$	\mathbf{on}
DUSU	ш		ULL

Reactivity Human, Mouse, Rat

Synthesized peptide derived from human Tak1 around the phosphorylation site of **Immunogen**

Thr187

Host Rabbit IgG **Isotype**

Purification Affinity purification

Conjugation **Unconjugated**

Formulation PBS with 0.02% sodium azide, 0.5% protective protein and 50% glycerol, pH7.4

Applications Recommended Dilution

WB 1:500-1:2000 **IHC** 1:100-1:300 1:10000 **ELISA**

Data



Western Blot analysis of 293T cells using Phospho-MAP3K7 (Thr187) Polyclonal Antibody at dilution of 1:1000

> Observed Mw:60kDa Calculated Mw:67kDa

Preparation & Storage

Storage Store at -20°C. Avoid freeze / thaw cycles.

Background

Component of a protein kinase signal transduction cascade. Mediator of TRAF6 and TGF-beta signal transduction. Activates IKBKB and MAPK8 in response to TRAF6 signaling. Stimulates NF-kappa-B activation and the p38 MAPK pathway. In osmotic stress signaling, plays a major role in the activation of MAPK8/JNK, but not that of NF-kappa-B.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com