

# Phospho-HDAC6 (Ser22) Polyclonal Antibody

Catalog Number:E-AB-21286



**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

|                     |   |
|---------------------|---|
| <b>Reactivity</b>   | Human,Mouse   |
| <b>Immunogen</b>    | Synthesized peptide derived from human HDAC6 around the phosphorylation site of Ser22 |
| <b>Host</b>         | Rabbit  |
| <b>Isotype</b>      | IgG   |
| <b>Purification</b> | Affinity purification   |
| <b>Conjugation</b>  | Unconjugated  |
| <b>Formulation</b>  | PBS with 0.02% sodium azide, 0.5% protective protein and 50% glycerol, pH7.4          |

## Applications Recommended Dilution

|              |              |
|--------------|--------------|
| <b>WB</b>    | 1:500-1:2000 |
| <b>IHC</b>   | 1:100-1:300  |
| <b>IF</b>    | 1:200-1:1000 |
| <b>ELISA</b> | 1:10000      |

## Data



Western Blot analysis of 3T3 cells with Phospho-HDAC6 (Ser22) Polyclonal Antibody at dilution of 1:500

**Observed Mw:131kDa**

**Calculated Mw:131kDa**

## Preparation & Storage

**Storage** Store at -20°C. Avoid freeze / thaw cycles.

## Background

Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Histone deacetylases act via the formation of large multiprotein complexes (By similarity). Plays a central role in microtubule-dependent cell motility via deacetylation of tubulin.

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017