

## APOBEC3A Polyclonal Antibody

**Catalog No.** E-AB-30528

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Reactivity</b>	Human
<b>Immunogen</b>	Synthesized peptide derived from the Internal region of human APOBEC3A
<b>Host</b>	Rabbit
<b>Isotype</b>	IgG
<b>Purification</b>	Affinity purification
<b>Conjugation</b>	Unconjugated
<b>Buffer</b>	PBS with 0.02% sodium azide, 0.5% protective protein and 50% glycerol, pH7.4

### Applications Recommended Dilution

<b>WB</b>	1:500-1:2000
<b>ELISA</b>	1:10000

### Data



Western Blot analysis of Jurkat cells with APOBEC3A  
Polyclonal Antibody.  
**Observed Mw:26kDa**  
**Calculated Mw:23kDa**

### Preparation & Storage

**Storage** Store at -20°C. Avoid freeze / thaw cycles.

### Background

This gene is a member of the cytidine deaminase gene family. It is one of seven related genes or pseudogenes found in a cluster, thought to result from gene duplication, on chromosome 22. Members of the cluster encode proteins that are structurally and functionally related to the C to U RNA-editing cytidine deaminase APOBEC1. The protein encoded by this gene lacks the zinc binding activity of other family members. The protein plays a role in immunity, by restricting transmission of foreign DNA such as viruses. One mechanism of foreign DNA restriction is deamination of foreign double-stranded DNA cytidines to uridines, which leads to DNA degradation. However, other mechanisms are also thought to be involved, as anti-viral effect is not dependent on deaminase activity. Two transcript variants encoding different isoforms have been found for this gene. APOBEC3A (Apolipoprotein B mRNA Editing Enzyme Catalytic Subunit 3A) is a Protein Coding gene. Among its related pathways are mRNA Editing- C to U Conversion and Gene Expression. GO annotations related to this gene include hydrolase activity, acting on carbon-nitrogen (but not peptide)

### For Research Use Only

bonds, in cyclic amidines and deoxycytidine deaminase activity. An important paralog of this gene is APOBEC3B.