CAMK2B/G Polyclonal Antibody

Catalog No. E-AB-30732

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human,Mouse,Rat
Immunogen	Synthesized peptide derived from the C-terminal region of human CaMKII β/γ
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Conjugation	Unconjugated
Buffer	PBS with 0.02% sodium azide, 0.5% protective protein and 50% glycerol, pH7.4
Applications	Recommended Dilution
WB	1:500-1:2000
IF	1:100-1:300
ELISA	1:20000
Data	



Western Blot analysis of COLO205 cells with CAMK2B/G Polyclonal Antibody Observed Mw:62kDa Calculated Mw:63kDa

Preparation & Storage

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Background

The multifunctional CAMK2, or Ca2+/calmodulin-dependent protein kinase II, is a well known effector of calcium- and calmodulin- mediated functions. It is present in many tissues but is most abundant in the brain. CAMK2 is composed of four different chains: alpha, beta, gamma, and delta. The different isoforms assemble into homo- or heteromultimeric holoenzymes composed of 8 to 12 subunits. Autophosphorylation plays an important role in the regulation of the kinase activity. CAMK2 is required for synaptic plasticity, as in Long Term Potentiation (LTP), a cellular model for learning and memory. CAMK2B is the beta chain. There are seven named isoforms produced by alternative splicing. The variable region of the CAMK2B protein is encoded by at least 7 exons (V1 to V7). Alternative splicing within this region gives rise to CAMK2B isoforms. CAMK2G is the gamma chain. CAMK2G has six alternatively spliced variants that encode six

For Research Use Only

Toll-free: 1-888-852-8623 Web: <u>www.elabscience.com</u>

Elabscience®

different isoforms. Some of these variants have been identified in human tumors.

For Research Use Only